

# Implementing 3-Year Protocols with Continuum™ DAP



**Continuum™**  
DAP

Better Medicine Through Customization™

“ I like the idea of 3-year vaccination with Continuum™ DAP. **BUT** how does it work with my other vaccines? ”

## Using the Sample Protocols

Simply decide if your patient's lifestyle and risk factors are most like Bailey, Tucker, King, or Coco. Then follow the corresponding vaccination protocol.

### BAILEY (high environmental exposure; high exposure to other dogs)

- The most common patient type
- Social; interacts with other dogs often
- Spends a lot of time outdoors
- Neighborhood lifestyle



| Year 1  | Year 2                       | Year 3                       | Year 4  | Year 5                       | Year 6                       | Year 7  | Year 8                       | Year 9                       |
|---|------------------------------|------------------------------|---|------------------------------|------------------------------|---|------------------------------|------------------------------|
| DAP<br>Rabies<br>Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme | DAP<br>Rabies<br>Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme | DAP<br>Rabies<br>Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme | Bb/PI/CAV-2<br>Lepto<br>Lyme |

### TUCKER (high environmental exposure; low exposure to other dogs)

- Rarely comes in contact with other dogs
- Spends a lot of time outdoors
- Farm dog lifestyle



| Year 1                         | Year 2        | Year 3        | Year 4                         | Year 5        | Year 6        | Year 7                         | Year 8        | Year 9        |
|--------------------------------|---------------|---------------|--------------------------------|---------------|---------------|--------------------------------|---------------|---------------|
| DAP<br>Rabies<br>Lepto<br>Lyme | Lepto<br>Lyme | Lepto<br>Lyme | DAP<br>Rabies<br>Lepto<br>Lyme | Lepto<br>Lyme | Lepto<br>Lyme | DAP<br>Rabies<br>Lepto<br>Lyme | Lepto<br>Lyme | Lepto<br>Lyme |

### KING (low environmental exposure; high exposure to other dogs)

- Around other dogs often
- Spends little time outdoors
- Show dog lifestyle



| Year 1                       | Year 2      | Year 3      | Year 4                       | Year 5      | Year 6      | Year 7                       | Year 8      | Year 9      |
|------------------------------|-------------|-------------|------------------------------|-------------|-------------|------------------------------|-------------|-------------|
| DAP<br>Rabies<br>Bb/PI/CAV-2 | Bb/PI/CAV-2 | Bb/PI/CAV-2 | DAP<br>Rabies<br>Bb/PI/CAV-2 | Bb/PI/CAV-2 | Bb/PI/CAV-2 | DAP<br>Rabies<br>Bb/PI/CAV-2 | Bb/PI/CAV-2 | Bb/PI/CAV-2 |

### COCO (low environmental exposure; low exposure to other dogs)

- Does not usually interact with other dogs
- Spends little to no time outdoors
- Pampered lifestyle



| Year 1        | Year 2 | Year 3 | Year 4        | Year 5 | Year 6 | Year 7        | Year 8 | Year 9 |
|---------------|--------|--------|---------------|--------|--------|---------------|--------|--------|
| DAP<br>Rabies |        |        | DAP<br>Rabies |        |        | DAP<br>Rabies |        |        |

DAP = Canine distemper virus, canine adenovirus-2, parvovirus  
Bb = *Bordetella bronchiseptica*  
PI = Canine parainfluenza

CAV-2 = Canine adenovirus-2  
Lepto = *Leptospira*  
Lyme = *Borrelia burgdorferi*

See back cover for information on vaccinating puppies with Continuum DAP.



# Vaccinating Puppies With a 3-Year Vaccine

You can safely and easily use Continuum™ DAP in even your youngest patients, as it can be administered to dogs as young as 6 weeks of age. Like most vaccines, Continuum DAP requires an initial vaccination and a booster vaccination. But after that, you simply vaccinate every 3 years.

The protocol below only shows core vaccines. Other vaccines should be incorporated into this protocol based on the patient's environmental and lifestyle risk factors. Also note that the protocol shows how 3-year Continuum DAP and a 3-year rabies vaccine can be staggered across different years.

## Puppy Protocol Guideline\*

| 8 weeks old | 12 weeks old  | 1 year old | 2 years old | 3 years old | 4 years old | 5 years old | 6 years old | 7 years old |
|-------------|---------------|------------|-------------|-------------|-------------|-------------|-------------|-------------|
| DAP         | DAP<br>Rabies | Rabies     |             | DAP         | Rabies      |             | DAP         | Rabies      |

\*Other vaccines (eg, *Borrelia burgdorferi* [Lyme], *Leptospira*, *Bordetella bronchiseptica*, etc) should be incorporated based on each patient's needs.



For information and assistance on incorporating Continuum DAP into your practice protocols, please contact your local distributor representative.

[www.continuum3.com](http://www.continuum3.com)

